

# Maximum Field Rodent Control

**rozol<sup>®</sup> RTU**

COMMERCIAL PRODUCT

- **Ready-To-Use:**  
*Premixed for labour savings*
- **Now approved for use on Pocket Gophers using tractor-driven burrow builder equipment**
- **Approved for preventative baiting in bait stations for Ground Squirrels**  
*Every ~100-200ft. (30-60m)*




20 lb. (9kg) pail

50 lb. (22.7kg) bag



Ground Squirrel



Learn More



Photo by Wayne Lynch

Pocket Gopher

Distributed in Canada by:  
**Univar Solutions 866.590.3444**

**rozol<sup>®</sup> RTU**

**Rids you of Ground Squirrels**

**LIPHATECH<sup>®</sup>**  
www.liphatech.ca

## UP TO C\$600 ROZOL RTU Rebate Offer UP TO C\$600

Redeem by mail - see reverse

<b>\$10<sup>00</sup></b> per bag (1-5 bags)	<b>\$12<sup>00</sup></b> per bag (6-15 bags)	<b>\$15<sup>00</sup></b> per bag (16-40 bags)
<b>\$4<sup>00</sup></b> per pail (1-5 pails)	<b>\$5<sup>00</sup></b> per pail (6-15 pails)	<b>\$6<sup>00</sup></b> per pail (16-100 pails)



**C\$600.00 max value**  
for up to 40 bags or 100 pails

For redemption questions, please call - (414) 351-1476 x7237

Limit one coupon per user/address, per 12 month period.

Offer valid for purchases from Feb. 10, 2020 - June 12, 2020.

Must be post marked by June 26, 2020.

Please allow 8-12 weeks for processing.

\* Rozol RTU (PCP #29545) is a commercial product. It is the responsibility of the user to read and follow all label directions.

Purpose: This rebate is for trial purchases by end-users only; not for redemption by resellers.

Offer valid in the provinces of: AB, BC, MB and SK.

Reproductions NOT accepted.

# rozol<sup>®</sup> MD RTU

## For Maximum Ground Squirrel & Pocket Gopher Control

Made with feed-grade winter wheat, this moisture-resistant formulation contains the anticoagulant chlorophacinone.

- **Season-long acceptance:** unlike strychnine that can spoil or may not be accepted by ground squirrels when other forage is available.
- **Moisture resistant:** can be used in wet or snowy conditions.
- **Easy-To-Use:** Rozol RTU is "Ready-To-Use". No more mixing or running out of grain or concentrate, or storage problems. Improve your cash flow and reduce potential waste by ordering only the bait you need when you need it.
- **Consistent blend:** no risk of under/over mixing of concentrate (active ingredient). Liphatech's rigorous quality control ensures that every kernel consumed contributes to effective rodent control.
- **Multiple use sites:** for use on rangelands, crop and non-crop areas to control ground squirrels.
- **Also controls pocket gophers** in alfalfa, hay and other forage crops, pulse/seed crops, cereal grains and on rangeland, non-crop areas, lawns and golf courses.
- **Preventative perimeter treatments intercept gophers** before they enter the field.
- **Vitamin K1 is the antidote.** There is no antidote for strychnine or zinc phosphide.
- Does **not** require use of protective eyewear or dust mask, as with strychnine.



Bait stations are an Effective Baiting Tool for Ground Squirrels

### Preventative Perimeter Treatments



*Only to be used by certified pest control operators, farmers, and persons authorized in government-approved pest control programs.*

## LIPHATECH<sup>®</sup>

For details, call Liphatech in Milwaukee, WI, at 1.888.331.7900, fax us at 414.247.8166, or visit us at [www.liphatech.ca](http://www.liphatech.ca)

Crops / areas where you experience highest incidence of damage?

Alfalfa  Cereal Grains  Legumes  Pulse/Seeds  Pasture List \_\_\_\_\_

Tell us what motivated you to try Rozol RTU? Is this your first time using Rozol RTU? (y/n)

\_\_\_\_\_

What was your experience with Rozol RTU? \_\_\_\_\_

Do you currently have at least four (4) bait stations per acre (on the perimeter) for a preventative baiting strategy? (y/n). Would you consider it to improve your results? (y/n). Why or Why not?

\_\_\_\_\_

I make burrow builder bait treatments after cuttings, whenever I see gopher mounds (y/n) \_\_\_\_\_

I make preventative burrow builder treatments around field borders \_\_\_\_ (y/n) \_\_\_\_ times/yr.

I apply bait (circle all that apply): Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov Dec

NAME: \_\_\_\_\_

ADDRESS: \_\_\_\_\_

CITY: \_\_\_\_\_ PROV: \_\_\_\_\_ POSTAL CODE: \_\_\_\_\_

PHONE: \_\_\_\_\_ EMAIL: \_\_\_\_\_

PARTICIPANT SIGNATURE\* \_\_\_\_\_

## rozol<sup>®</sup> MD RTU Rebate Offer

### Instructions:

Mail completed coupon along with a dated purchase receipt to:

Univar Agriculture  
Rozol RTU Rebate  
99 Lowson Crescent  
Winnipeg MB R3P 0T3  
Canada

*Incomplete surveys may be denied payment.*

*\*Authorizing Liphatech limited use of comments in printed promotional materials and publicity-related communications. Your name will NOT be used or SOLD for any other purpose.*

COMMERCIAL

# ROZOL® RTU

## Field Rodent Bait Ready-To-Use

Only to be used by certified pest control operators, farmers and persons authorized in government-approved pest control programs.

WARNING /  
POISON



ACTIVE INGREDIENT: Chlorophacinone..... 0.005%  
REGISTRATION NO. 29545 PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN  
READ THE LABEL BEFORE USING  
WARNING - CONTAINS THE ALLERGEN WHEAT

**LIPHATECH®**

Liphatech, Inc.  
3600 W. Elm Street  
Milwaukee, WI 53209 USA  
(800) 351-1476

**USE LIMITATIONS - GENERAL:** DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**DIRECTIONS FOR USE:** Rozol® RTU Field Rodent Bait may be used for the control of Pocket Gophers, *(Thomomys spp. and Geomys spp.)*, In lawns, golf courses, alfalfa, hay and other forage crops, cereal grains, range land, and non-crop areas and Ground Squirrels (*Citellus richardsoni*, *U. columbianus*, *Poleciolus flaviventris*, *Citellus tridecemlineatus*) in pulse-seed crops, rangelands, crop and non-crop areas.

Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing. Replace contaminated or soiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water and harborage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as necessary.

**GROUND SQUIRRELS** (Richardson, Columbian, Franklin's and Thirteen-lined):

**Areas of Use - Ground Squirrels:** Ground squirrels typically inhabit range lands, pulse/seed crops and areas directly adjacent to crops. Even though ground squirrels are foraging animals, placement of baits directly into the burrows or the use of bait stations is required to reduce hazards to children and non-target animals.

**Application Directions - Ground Squirrels - Hand Baiting:** Locate at the ground squirrel burrows in the area of treatment and place 7 to 14 grams (0.25 to 0.50 ounces) of bait into each burrow. Re-visit the site within 48 hours and re-bait all burrows where bait has been consumed or removed. A third re-baiting visit may be necessary for control. Do not re-bait burrows where the bait is undisturbed. Remove and bury any squirrel carcasses which are present above ground.

**Bait Station Baiting:** Determine the approximate population present by visual count over a two-day period. In areas where ground squirrels are present, establish protected bait stations or tamper resistant boxes with an entrance hole of 7.6 cm (3 inches) using 1 station for every 50 animals counted. Place 500 grams (17.6 ounces) of bait into the stations and check daily. Maintain an uninterrupted supply for 3 weeks or until all signs of feeding have stopped. In areas where ground squirrels have been known to invade, but are not currently present, preventative baiting in stations can be accomplished by placing stations containing 500 grams (17.6 ounces) of bait every 30 to 60 metres (100 to 200 feet). Replace spoiled or moldy bait as necessary. Remove and bury any squirrel carcasses which are present above ground.

**USE LIMITATIONS - Ground Squirrels:**

Bait placed outdoors and above-ground in residential areas, public areas, or any other sites that are or will be open to the public (for example, Pet-Your-Own sites) or in locations accessible to pets or livestock MUST be placed in Tier 1 bait stations.

Tier 1 bait stations must have the following characteristics:

1. constructed of high-strength material (e.g., metal or injection-moulded plastic) and resistant to destruction by children and non-target animals;
2. have an entrance designed so that children and non-target animals cannot reach the bait;
3. have an internal structure that prevents bait from being stolen loose;
4. have an access panel which lasts securely and locks (e.g., metal screw or padlock);
5. bear the product name, active ingredient, guarantee, registration number, "WARNING POISON", and the skull and crossbones symbol;
6. be resistant to destruction or weakening by elements of typical non-catastrophic weather (such as snow, rain, extremes of temperature and humidity, direct sunshine, etc.).


**POCKET GOPHERS:** *(Thomomys spp. and Geomys spp.)*:

**Areas of Use - Pocket Gophers:** Pocket gophers throw out low, fan-shaped mounds on either side of their underground burrows. These lateral tunnels come to the surface and are on the concave or flat side of the fan, and the entrance holes are plugged with loose soil.

**Application Directions - Pocket Gophers:**

Placement of the bait can be made in any of the following ways:

1. **With a long-handled spoon,** carefully remove the plug on the flat side of the fan. Carefully insert 15 grams (0.53 ounces) of bait as far down into the hole as possible. Replace the soil plug to close the burrow, with care so that the bait is NOT covered with soil. Make 2-3 bait placements per burrow system. Bury dead animals and spoiled bait found on soil surface immediately. Maintain a constant supply of bait in the burrow system for as long as there is pocket gopher activity. Do not apply bait on the surface of soil.
2. **Using a metal rod,** probe 15 to 20 centimetres (6 to 8 inches) deep to locate the main tunnel. Consult diagram below for location to probe. Drop 15 grams (0.53 ounces) of bait into the tunnel and cover the hole so light will not enter the tunnel system.



The right and the wrong ways to use a probe for placing pocket gopher bait are shown. Bait should be placed loosely in the main runway - not in the lateral tunnels or embedded in the bottom of the runway.

Make 2-3 bait placements per burrow system. Bury dead animals and spoiled bait found on soil surface immediately. Maintain a constant supply of bait in the burrow system for as long as there is pocket gopher activity. Do not apply bait on the surface of soil.

**3. Machine application using "burrow builder" equipment:** Follow manufacturer's instructions for the equipment used to apply this bait. Make artificial burrows at the same depth as natural burrows, about 4.8 to 9 metres (15 to 30 feet) apart.

Apply 1.7 to 4.5 kilograms of bait per hectare (1.5 to 4 pounds per acre). Use the high rate in locations with greater pocket gopher activity. Apply when soil conditions allow formation of good artificial burrows. Do not treat the entire field. Apply as:

1. A targeted treatment to the part of the field where pocket gophers have established burrows. Make 6 to 8 rows in the area infested with pocket gophers. Do not exceed 3 applications per year.
2. Perimeter treatment to reduce potential migration of pocket gophers into the field. Make 3 rows around the perimeter of the field. Use only in locations where there is a risk of pocket gopher damage. Do not exceed 3 applications per year; one each in the fall, spring and mid-summer.

**PRECAUTIONS - KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.** May be harmful or fatal if swallowed or absorbed through the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents. Unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODS UPFS.** All handlers must wear long-sleeved shirt and long pants, shoes plus socks, and chemical-resistant gloves when handling this product.

**ENVIRONMENTAL PRECAUTIONS:** Toxic to aquatic organisms, birds and small mammals. Use only in sites listed on the product label. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

**FIRST AID - IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. Call a poison control centre or doctor for further treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.

For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin K<sub>1</sub> in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**STORAGE AND DISPOSAL:** Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original, tightly sealed container, in a secure location inaccessible to children and non-target animals.

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

**CARCASS DISPOSAL:** To protect pets and non-target wildlife, carefully dispose of animals killed by this bait. Wrap poisoned carcasses securely with paper. Dispose of them by placing in a closed container and taking them to a provincial or municipal landfill site OR burying deep enough to prevent scavengers from reaching them.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. (112218)

