DIRECTIONS FOR USE:

Use DoubleTap Soft Bait for the control of Norway rats (*Rattus norvegicus*), Roof rats (*Rattus rattus*), House mice (*Mus musculus*), Deer mice (*Peromyscus maniculatus*) and White-footed mice (*P. leucopus*) indoors and outdoors within 15 metres of buildings (residential, industrial, commercial, agricultural, and public buildings). Rodenticide bait can be placed in tamper-resistant bait stations along the fence line of properties, outside of the 15-metre limit but within 100 metres of buildings, if the station is securely fastened (e.g., nailed down) to the fence or the ground. This product may also be applied in garbage dumps.

Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing.

USE LIMITATIONS: Bait MUST be placed either in tamper-resistant bait stations or in locations not accessible to children, pets, livestock or non-target wildlife. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

Bait stations (tier 3) used for the placement of rodenticide bait indoors, in locations not accessible to pets or livestock must have the following characteristics: (1) be constructed of high-strength material (e.g., metal or injection moulded plastic) and resistant to destruction by children; (2) have an entrance designed so that children cannot reach the bait; (3) have an internal structure that prevents bait from being shaken loose; (4) have an access panel that fastens securely and locks (e.g., metal screw or padlock); and (5) bear the product name, active ingredient, guarantee, registration number, "WARNING POISON", and the skull and crossbones symbol.

Bait stations (tier 2) used for the placement of rodenticide bait indoors, in locations accessible to pets or livestock must have the following characteristics, in addition to those outlined above for tier 3 bait stations: (1) resistant to destruction by non-target animals; and (2) have an entrance designed so that non-target animals cannot reach the bait.

Bait applied outdoors and above-ground MUST be placed in bait stations. Bait stations (tier 1) used outdoors, above-ground, in locations accessible to children, pets and non-target wildlife must have the following characteristic, in addition to those outlined above for tier 2 and 3 bait stations: (1) be resistant to destruction or weakening by elements of typical non-catastrophic weather (such as. snow, rain, extremes of temperature and humidity, direct sunshine, etc.).

Food Processing, Food Manufacturing, Food Storage and Food Service Areas: For areas not directly related to food processing: Use only in non-food or non-feed areas where feed, food, packaging and handling equipment are never opened or exposed. For areas where feed or food is processed, served or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents and wash treated areas before re-use of the food handling establishment (processing areas include storage and service).

Selection of Treatment Areas: Place bait in locations where listed rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and near burrows, and in concealed locations where listed rats and mice feed is desirable. Eliminate as much food, water and harborage as possible. Use the highest rate permitted on this label in areas of very high pest activity.

APPLICATION DIRECTIONS: For Norway rats and Roof rats: Norway rats usually inhabit the ground floor of structures and burrow in the ground. They will range approximately 15 to 100 m from their nesting area seeking food and water. Roof rats are excellent climbers and usually infest the upper stories of structures. Hence, special attention must be paid to this species ability to climb by placing baits in the roof rats' activity area. Apply 6 to 16 pouches per placement, usually spaced 5 to 10 m apart. Maintain a constant supply of fresh bait for 10 days or until signs of rat activity cease.

For House mice, Deer mice and White-footed mice: House mice, dear mice and white-footed mice are excellent climbers and will infest all floors in a structure. House mice will only range approximately 10 m from their nesting area seeking food. Deer mice and white-footed mice will travel further in search of food. If a bait placement is not disturbed after several days, move the placement approximately 1 metre. Apply 4 to 6 pouches per placement, usually spaced 2 to 4 m apart. Maintain a constant supply of fresh bait for 15 days or until signs of activity cease.

Note: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait as described under DISPOSAL. To prevent reinfestation, eliminate rodent food, water and harbourage as much as possible. If reinfestation occurs, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.



RODENTICIDE

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE, DEER MICE, AND WHITE-FOOTED MICE

PASTE COMMERCIAL

ACTIVE INGREDIENT: Chlorophacinone...... 0.005%

WARNING, CONTAINS THE ALLERGENS WHEAT AND SOY

REGISTRATION NO. 34220

PEST CONTROL PRODUCTS ACT

KEEP OUT OF REACH OF CHILDREN READ THE LABEL BEFORE USING

Only to be used by certified pest control operators, farmers and persons authorized in government-approved pest control programs.



Liphatech, Inc. 3600 W. Elm Street Milwaukee, WI 53209 USA (800) 351-1476 PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed through the skin. **Do not** open premeasured place packs (pouches). Wear a long-sleeved shirt, long pants, chemicalresistant gloves, socks and shoes when handling this product, disposing of dead
rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or
clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using
the toilet. Wash skin thoroughly with soap and water after handling. Users should
remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and
put on clean clothing. Wash contaminated clothing, separately from other laundry,
with soap and hot water before reuse. Do not place bait in areas where there is a
possibility of contaminating food or surfaces that come in direct contact with food.
KEEP AWAY FROM FEED AND FOODSTUFFS

centre or doctor.	FIRST AID:	
Rinse skin with plenty of water for 15-20 minutes. Call a poison control centre or doctor immediately for treatment advice. If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce womiting unless told to do so by a poison control centre or doctor.	If in eyes:	15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor immediately for
treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomitting unless told to do so by a poison control centre or doctor.		Rinse skin with plenty of water for 15-20 minutes. Call a poison control centre or doctor immediately for
For all cases of human ingestion call a poison control centre or doctor		treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control centre or doctor. • Do not give anything by mouth to an unconscious person.

immediately for treatment advice.

If pet or livestock poisoning is suspected, immediately contact a veterinarian.

If pet or livestock poisoning is suspected, immediately contact a veterinarian.

Take container, label or product name and Pest Control Product Registration

Number with you when seeking medical attention.

 $\label{thm:colored} \textbf{TOXICOLOGICAL INFORMATION:} \quad \text{This material may reduce the clotting ability of blood and cause bleeding. Vitamin <math>K_1$ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

ENVIRONMENTAL PRECAUTIONS: Toxic to aquatic organisms, birds and small wild mammals. Use only in sites listed on the product label. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

STORAGE: Store this product away from food or feed. Store product not in use in original, tightly sealed container in a cool, dry, secure location away from other chemicals and inaccessible to children and non-target animals.

DISPOSAL:

CONTAINER: Do not reuse empty container. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

PRODUCT: For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

CARCASSES: To protect pets and non-target wildlife, carefully dispose of animals killed by this bait. Wrap poisoned carcasses securely with paper. Dispose of them by placing in a closed container and taking them to a provincial or municipal landfill site OR burying deep enough to prevent scavengers from reaching them.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. (083121)